

Submitted to *The Astrophysical Journal* in final form 8 May 1999

No Hubble Bubble in the Local Universe

Riccardo Giovanelli, Daniel A. Dale, Martha P. Haynes

Center for Radiophysics and Space Research and National Astronomy and Ionosphere Center, Cornell University, Ithaca, NY 14853

Eduardo Hardy

National Radio Astronomy Observatory, Casilla 36-D, Santiago, Chile

Luis E. Campusano

Observatorio de Cerro Calán, Universidad de Chile, Casilla 36-D, Santiago, Chile

ABSTRACT

Zehavi et al. (1998) have suggested that the Hubble flow within $70h^{-1}$ Mpc may be accelerated by the existence of a void centered on the Local Group. Its underdensity would be $\sim 20\%$, which would result in a local Hubble distortion of about 6.5% . We have combined the peculiar velocity data of two samples of clusters of galaxies, SCI and SCII, to investigate the amplitude of Hubble distortions to $200h^{-1}$ Mpc. Our results are not supportive of that conclusion. The amplitude of a possible distortion in the Hubble flow within $70h^{-1}$ Mpc in the SCI+SCII merged data is 0.010 ± 0.022 . The largest, and still quite marginal, geocentric deviation from smooth Hubble flow consistent with that data set is a shell with $\Delta H_o/H_o = 0.027 \pm 0.023$, centered at $hd = 101$ Mpc and extending over some $30 h^{-1}$ Mpc. Our results are thus consistent with a Hubble flow that, on distances in excess of $\sim 50h^{-1}$ Mpc, is remarkably smooth.

Subject headings: galaxies: distances and redshifts – cosmology: observations; cosmic microwave background; distance scale

1. Introduction

The linearity of the Hubble law over large scales, as illustrated by the early work of Sandage & Hardy (1973), has been confirmed by more recent measurements as discussed by Postman (1997). These measurements do not however exclude the possibility of local deviations from Hubble flow with amplitudes on the order of a few percent, as would be produced by large-scale mass fluctuations. For example, an extended, underdense region centered on the Local Group would exhibit a locally accelerated Hubble flow. This could, to a point, help reconcile discrepant estimates of the value of the Hubble constant obtained by methods which sample vastly different scales and solve the still raspy conflict between some estimates of the age of a matter dominated Universe and that of the oldest stars. By analyzing the monopole of the peculiar velocity field as described by a sample of 44 type Ia supernovae (SN), Zehavi *et al.* (1998, hereafter Z98) have recently suggested that the Local Group may be near the center of a bubble of $70h^{-1}$ Mpc radius (where $H_o = 100h$ km s $^{-1}$ is the Hubble constant), underdense by 20% , which may be itself surrounded by an overdense shell. The isotropic flow observed within that “bubble” would then exceed the universal rate by $\Delta H_o/H_o = (6.5 \pm 2.1)\%$; i.e., studies that rely on distance indicators contained within that bubble

would overestimate the Hubble constant by 6.5% . Z98 cautiously underscore the marginal character of their detection, as well as the need to corroborate, or refute, their suggestion by means of tests with independent sets of data. In this *Letter*, we provide such a test.

2. The SCI and SCII Cluster Samples

Based on data published earlier (Giovanelli *et al.* 1997a,b; hereafter G97a and G97b), peculiar velocities of 24 clusters of galaxies within $90h^{-1}$ Mpc, obtained from measurements for 782 galaxies in their fields (hereafter referred to as SCI) have been recently presented by Giovanelli *et al.* (1998a), who also used it to compute a dipole and investigate the Z98 claim. Because of the limited depth of the SCI sample, their test of the Z98 claim was inconclusive. Recently, we have completed a deeper survey of cluster peculiar velocities, which extends to $200h^{-1}$ Mpc. As for the SCI sample, the new survey is based on the Tully–Fisher (1977; hereafter TF) technique. The first installments of this data set are in Dale *et al.* (1997, 1998); the final one is in preparation, but in preliminary form its results can be seen in Dale (1998). The new survey, which we shall refer to as SCII, includes 522 galaxies in 52 clusters. The dipole signature of the SCII, which is consistent with that of the CMB temperature dipole, is discussed in Dale *et al.* (1999).

The combination of SCI and SCII provides a peculiar velocity data set of slightly smaller depth, but higher sampling density than the SN sample of Z98. The peculiar velocity errors of the SCI set vary between 3% and 6% of the distance, for each individual cluster. In the case of the SCII set, peculiar velocity errors are somewhat higher, due to the smaller number of galaxies observed per cluster: they hover between 4% and 9%, except in a few cases which will be discussed later. On the average, the accuracy of each cluster peculiar velocity compares favorably with the quoted uncertainty of 5–8% (for the internal errors alone) of the distance of individual SN in the Z98 sample. Since the 76 clusters in the SCI+SCII merged sample straddle quite comfortably the boundaries of the Z98 bubble, they can provide tighter constraints than the SN sample on the amplitude of the proposed, locally underdense region.

Table 1 lists the clusters in the SCI+SCII merged sample, identified either by their Abell number (Abell, Corwin & Olowin 1989) or by their common name, the adopted center coordinates, as well as the radial velocity cz_{cmb} and the peculiar velocity V_{pec} in the CMB reference frame, (after Giovanelli *et al.* 1998a and Dale 1998) and the number N of galaxies in each cluster with TF measurements. We compute a distance $hd = (cz_{cmb} - V_{pec})/100$ and a deviation from Hubble flow $\Delta H_o/H_o = V_{pec}/(cz_{cmb} - V_{pec})$.

The total error on the peculiar velocity of each cluster, as listed in Table 1, includes several components, arising from: (i) photometric and spectroscopic observational errors; (ii) uncertainties in the corrections applied to observed parameters; (iii) uncertainties in the cluster redshifts; (iv) the scatter in the TF relation; (v) uncertainties in the TF template relation slope and zero point, especially that deriving from the assumed standard of rest. We discuss point (v) in greater detail in the next Section. The other sources of error are extensively discussed in the data papers mentioned above.

3. Template Relation Accuracy and its Effect on the Monopole Moment

TF peculiar velocities are derived as offsets from a template relation, which in its simplest form is defined by two parameters: a slope and a zero point. Errors on both the zero point and on the slope translate into spurious, geocentric peculiar velocity fields. For example, an error of 0.05 mag in the zero

point would simulate a slowdown or speeding up of the Hubble expansion by 2.3%. As for the effect of an error on the TF slope: if the template relation is, for example, too steep — i.e. for a given velocity width which is broader than some fiducial value the template predicts too bright a magnitude —, then high width galaxies will preferentially yield positive magnitude offsets. The opposite will be true for low width galaxies. Since low width galaxies are intrinsically faint, they are more likely to be present in nearby samples than in more distant ones; thus nearby samples fitted with too steep a TF template relation exhibit a net negative magnitude offset, which translates into a spurious outflow. The effect of unrecognized TF calibration errors can then be misconstrued as a monopole perturbation, and thus as a geocentric Hubble flow distortion.

The TF template relation is determined *internally* for a cluster sample. In the case of SCI, it was obtained by assuming that the subset of clusters farther than $40h^{-1}$ Mpc has a globally null monopole (G97b). Dale (1998) obtained an SCII template by assuming that the set of clusters has a globally null monopole, and adopting the same TF slope as for the SCI sample. As discussed in G97b, given a number N of clusters the uncertainty on the TF zero point of the resulting template cannot be depressed indefinitely by increasing the average number $\bar{n}$ of galaxies observed per cluster, and taking advantage of the $\bar{n}^{-1/2}$ statistical reduction of noise on the mean. That is because a “kinematical” or “thermal” component of the uncertainty depends on the number N , distribution in the sky and peculiar velocity distribution function of the clusters used. In SCI, for example, the statistical uncertainty deriving from the total number of galaxies observed ($\bar{n} \times N$) is exceeded by the kinematic uncertainty, which is quantified as follows. For a sample of N clusters of average redshift $\langle cz \rangle$, the most probable systematic error on the template relation zero point is $|\Delta m| \simeq 2.17 \langle V_{pec}^2 \rangle^{1/2} \langle cz \rangle^{-1} N^{-1/2}$, where $\langle V_{pec}^2 \rangle^{1/2}$ (expressed in the same units as cz) is the line of sight r.m.s. cluster peculiar velocity, of about 300 km s^{-1} (G97b; Giovanelli *et al.* 1998a; Dale 1998). This quantity is about 0.04 mag for SCI, while it is only 0.01 mag for SCII due to the larger mean distance and number of clusters of the latter. Since the total number of galaxies involved in the two samples is comparable, the zero point of the SCII template is thus more accurate than that of SCI. On the other hand, the peculiar velocities of individual clusters in SCII are less accurate than those in SCI. We note that the kinematical or thermal component of the uncertainty is larger for SN peculiar velocities than for our cluster ones. That is because the amplitude of the distribution function of peculiar velocities among individual galaxies — the hosts of SN — is larger than that of clusters, as the former is amplified by the variance associated with fluctuations on small scales.

In the case of both SCI and SCII, a direct TF template relation was obtained, using the approach described in G97b. The data for each cluster offset was corrected for the effect of an incompleteness bias. The zero points of the two templates were found to agree to within 0.015 mag (SCII being fainter by that amount).

In this paper, we combine the SCI and SCII samples, and use them to investigate the presence of large scale variations in the monopole of the Hubble flow. Note that such combined sample cannot be used for the detection of a geocentric deviation from smooth Hubble flow *which would extend over the full volume sampled by the total cluster set, as it would be null by design*. The merged cluster data set can however be used to detect *changes in $\Delta H_o/H_o$* that would take place well within the volume spanned by the data. The amplitude of the change (say a step in $\Delta H_o/H_o$) that can be detected depends on the location of the presumed step and on the accuracy with which the match in the TF zero point between the SCI and SCII samples is established:

- (i) For our cluster sample, a step would be ideally situated between $70h^{-1}$ and $110h^{-1}$ Mpc, in order to maximize the chance of detection, because it would split the cluster sample into two roughly equal parts. The SCI+SCII sample is thus well-suited to test the Z98 result.

(ii) The internal accuracy of the zero point for the SCI sample is 0.025 mag; however, since it is based on a subset of 14 clusters farther than $40h^{-1}$ Mpc, the kinematical uncertainty of 0.04 mag, as mentioned above, increases the total uncertainty to 0.045 mag. The total uncertainty on the zero point of SCII, because it involves a larger number of more distant clusters, is only 0.025 mag; the kinematical component in this case is only 0.01 mag. Note for comparison that a 6.5% step in $\Delta H_o/H_o$ would translate in a 0.13 mag differential TF offset between clusters on each side of the step.

It is also useful to point out that each of the two samples was completed over many observing runs, both in their photometric and spectroscopic parts, and a number of objects were observed in more than one run. Mismatches in the cross-run and cross-cluster calibrations thus have been minimized and their impact on the final error budget is included in the statistical estimate given above.

(iii) The small overlap in distance between SCI and SCII occurs near $70h^{-1}$ Mpc, which is the edge of the Hubble bubble suggested by Z98 (4 clusters in SCI are farther than $cz = 7000 \text{ km s}^{-1}$, while 4 in SCII are within that redshift). We thus need to establish the impact of the accuracy of the match between the two samples' zero points, on the estimation of the likelihood of a Hubble bubble. We return to this point in Section 4.1.

4. Geocentric Hubble Deviations

Using the data in Table 1, and forcing the template TF zero point to be the same for SCI and SCII, we obtain Figure 1, a plot of the Hubble deviation versus the distance. In the upper panel of Figure 1 we display the individual data points, while in the lower one we show the errors associated with each measurement. Starred symbols refer to the SCI sample, while circular symbols refer to SCII. Eight clusters, flagged by double daggers in Table 1, are plotted in Figure 1 as unfilled symbols: their peculiar velocities have been obtained from fewer than five TF measurements and are thus very unreliable. The latter are not used in the following statistical analyses.

The plot presented in Figure 1 is similar to that in Figure 1 of Z98. For comparison, we have included the outline of the Z98 step as a dashed line, which extends from zero to $70h^{-1}$ Mpc distance, at the level of $\Delta H_o/H_o = 0.065$. We note immediately that the Z98 proposal of a Hubble bubble is not corroborated by the cluster data. We also note that at distances nearer than $\sim 30h^{-1}$ Mpc even modest peculiar velocities constitute a sizable fraction of cz , thus amplifying and distorting the values of $\Delta H_o/H_o$. The implied deviation from Hubble flow they reveal is of scarce interest, as they apply to too small, too sparsely sampled a volume.

Next, we test for the presence of a step at $70h^{-1}$ Mpc distance, of the kind suggested by Z98, and we search for the signature of other possible, geocentric large-scale fluctuations in the Hubble flow.

4.1. Test for a Hubble Bubble

We consider whether a step is present in $\Delta H_o/H_o$ at $70h^{-1}$ Mpc, by taking the difference in the average of $\Delta H_o/H_o$ between $30h^{-1}$ and $70h^{-1}$ Mpc, and the corresponding average at distances higher than $70h^{-1}$ Mpc. That difference is 0.010 ± 0.012 , if individual clusters are weighed by their errors in $\Delta H_o/H_o$, and 0.007 ± 0.012 if equal weight averages are computed. The uncertainty of this result can however be affected by a number of systematic errors, which exceed the statistical estimate given above; in the following we

discuss them one by one.

Kinematic Zero Point Mismatch: First, we consider the impact of the systematic mismatch between the TF template zero points of the two samples, as discussed in part (iii) of Section 3. We can evaluate the impact of that uncertainty on the determination of the amplitude of a possible step at $70h^{-1}$ Mpc by offsetting by $\pm(0.04^2 + 0.01^2)^{1/2} = \pm 0.04$ mag the SCI and SCII samples with respect to each other, computing in each case the amplitude of the step (note that there are SCI and SCII clusters on both sides of the step). The results are respectively 0.022 and -0.004 mag. It can thus be inferred that the impact on the uncertainty of the step, produced by a possible systematic error in the match between zero points for the two samples, is about 0.03 mag or 1.5%.

Differential Malmquist Bias: Malmquist bias corrections have not been applied to the cluster peculiar velocities. If such a correction were the same for all the clusters, it would have no impact on the detectability of a Hubble bubble step. However, since the more distant clusters of SCII each include a smaller number of galaxies with TF measurements, the impact of a possible differential Malmquist bias between SCI and SCII needs to be explored. As discussed in Giovanelli *et al.* (1998a), the Malmquist bias can be estimated with adequate accuracy in the "homogeneous" assumption, i.e. that the clusters' distribution in space is Poissonian, and shown to be quite small. The Malmquist bias correction in that case is $e^{3.5\Delta^2} - 1$, where $\Delta = 10^{0.2\epsilon/\sqrt{n}} - 1$, with ϵ the scatter in magnitudes about the TF relation (about 0.35 mag) and n the number of galaxies with TF measurements per cluster. For example, for a cluster with 10 galaxies with TF measurements, the average for SCII, the Malmquist bias correction is 1.0% on the distance. In the case of SCI, the average number of galaxies with TF measurements per cluster is about 16. In that case the Malmquist bias correction is 0.7% on the distance. Neglect to apply a Malmquist bias correction thus introduces a possible bias with an amplitude of 0.003 in $\Delta H_o/H_o$.

Template Relation Slope: The same template relation slope has been used for both SCI and SCII, as discussed in Dale (1998). The error on the determination of that slope is given in G97b, as 0.12 on a slope of -7.68, or 1.6%. If there were a significant difference in the distribution of galaxies as a function of velocity width, between nearby and more distant clusters, the uncertainty on the slope would introduce a systematic bias in the distances. To estimate the amplitude of that bias, we binned galaxies as a function of width, separately for the clusters within and beyond $70h^{-1}$ Mpc, and for each group estimated the average magnitude offset introduced by an error in the slope of 1.6%; in doing so, we assumed that the zero point, i.e. the value of the template relation at $\log W = 2.5$, is correct. The resulting TF offset uncertainty between the two groups is 0.0055 mag, or 0.0025 on the distance.

Evolution: Some authors (Rix *et al.* 1997; Simard & Pritchett 1998) have claimed substantial evolution in the mass-to-light ratio of spiral galaxies between $z = 0$ and relatively modest redshifts $z \sim 0.4$, while others (Vogt *et al.* 1997; Bershadsky 1996; Dale, Usón & Giovanelli 1999, in preparation) find no such effect. Evolution would translate into a shift of the TF relation zero point. While this issue is still quite uncertain, we can estimate the possible impact of evolution, assuming a (rather generous) shift of 1 mag between $z = 1$ and $z = 0$. The difference in z between the clusters within $70h^{-1}$ Mpc and those farther away is $\simeq 0.02$, thus a possible shift of 0.02 mag or 0.01 in distance would be possible. The direction of this relative shift would be that of a gradual brightening of the higher redshift galaxies and therefore increasing their average $\Delta H_o/H_o$. Probably overestimating it, we conclude that the uncertainty associated with this effect is 0.01 in the distance.

Below, we give a summary of the components of uncertainty with which the SCI+SCII merged sample can be used to identify a possible step in the Hubble flow at $70h^{-1}$ Mpc:

statistical distance error	1.2%
kinematic zero point mismatch between SCI and SCII	1.5%
Malmquist bias	0.3%
Template slope uncertainty	0.3%
Evolution	1.0%
Total	2.2%

Thus our estimate of the amplitude and significance of a step in the Hubble flow at $70h^{-1}$ Mpc is

$$\frac{\Delta H_o}{H_o} = 0.010 \pm 0.022 \quad (1)$$

For a two-zone model, which includes an inner void out to $70h^{-1}$ Mpc and an outer region expanding at the Hubble rate, Z98 report an amplitude of 0.065 ± 0.021 ; such a void is not apparent in our data, in which a 6.5% step would be a $3\text{-}\sigma$ event. We remark however that, compounding our estimated error with that reported in Z98, the difference between the two results is 0.055 ± 0.030 .

The Hubble distortion reported by Z98 reduces to 0.053 ± 0.022 in a three-zone model, where an inner underdense sphere of $70h^{-1}$ Mpc is surrounded by an overdense shell between 70 and $105 h^{-1}$ Mpc; for the outer shell, Z98 report an inflow of $\Delta H_o/H_o = -0.059 \pm 0.027$. For the latter region, between 70 and $105 h^{-1}$ Mpc, our data yields $\Delta H_o/H_o = +0.020 \pm 0.018$. The difference between our and the Z98 results, compounding the reported errors, is 0.079 ± 0.032 .

4.2. Other Geocentric Deviations

We search for the signature of other possible large-scale fluctuations in the Hubble flow, adopting a similar χ^2 minimization analysis to that carried out by Z98. The minimization of

$$\chi^2 = \sum_i \left[\log[1 + (\Delta H_o/H_o)_i] - \log[1 + (\Delta H_o/H_o)_{model}] \right]^2 / \sigma_i, \quad (2)$$

where $(\Delta H_o/H_o)_i$ are the values plotted in Figure 1 and $(\Delta H_o/H_o)_{model}$ is a model with a constant departure from zero in $\Delta H_o/H_o$ between two arbitrary distances hd_1 and hd_2 , is equivalent to that expressed in Eqn. (2) of Z98. σ_i is the estimated error on $\log[1 + (\Delta H_o/H_o)_i]$. The strongest signature for a departure from Hubble flow consistent with the SCI+SCII cluster data is a (very marginal) step of amplitude $\Delta H_o/H_o = 0.027 \pm 0.023$ centered at $hd = 101$ Mpc and 33 Mpc wide. The boundaries of the region are very “soft”. In the calculations, we impose that the width of the perturbed region should be no less than 20 Mpc, including only 61 clusters with hd between 30 and 200 Mpc and excluding the eight clusters with poor sampling ($n < 5$) plotted as unfilled symbols in Figure 1 and flagged in Table 1.

5. Conclusions

Z98 have cogently argued that a region of $70h^{-1}$ Mpc radius could be underdense by $\sim 20\%$ — which is the amount necessary to produce a suggested local acceleration of 6.5% of the Hubble flow —, without unreasonably stretching the plausible amplitude range of cosmological density fluctuations. One would be left, of course, with the nagging coincidence of the nearly central location of the LG in the void (a circumstance which would also be at some odds with the fairly large peculiar velocity of the LG of some 620




Fig. 1.— Deviations from Hubble flow plotted versus TF distance for the clusters listed in Table 1. In the upper panel, starred symbols refer to clusters in SCI, while circular symbols refer to SCII. Among the latter, filled symbols identify clusters with distance determinations based on $n > 4$ individual galaxy TF distances, while unfilled ones refer to clusters with $n \leq 4$, the peculiar velocities of which are deemed least trustworthy and are not used in the statistical analysis; their names are flagged in Table 1. The horizontal dashed line identifies the acceleration of 6.5% in the Hubble flow within $hd = 70$ Mpc claimed by Z98. The lower panel displays the error bars associated with each starred or filled data point.

km s⁻¹, as indicated by the CMB dipole). Our data give an amplitude for a possible Hubble flow distortion within 70 h^{-1} Mpc of $\Delta H_o/H_o = 0.010 \pm 0.022$.

In a three-zone model, Z98 suggest that an overdense shell between 70⁻¹ and 105 h^{-1} Mpc may be affected by an inflow of $\Delta H_o/H_o = -0.059 \pm 0.027$. For that region, our data yields $\Delta H_o/H_o = 0.020 \pm 0.018$.

The distortion of largest amplitude, consistent with our data, is $\Delta H_o/H_o = 0.027 \pm 0.023$ centered at $hd = 101$ Mpc and extending over a shell some 30 h^{-1} Mpc thick.

The results of this paper are consistent with those on the peculiar velocity field as traced by the SFI sample of field spirals: its dipole converges to that of the CMB dipole, both in amplitude and apex direction, within about 50 h^{-1} Mpc (Giovanelli *et al.* 1998b). We conclude that, at distances in excess of $\sim 50h^{-1}$ Mpc, the cluster peculiar velocity data are consistent with a picture in which the average Hubble flow is remarkably smooth.

The comments of an anonymous referee helped improve the presentation of this paper. The results presented in this paper are based on observations carried out at the Arecibo Observatory, which is part of the National Astronomy and Ionosphere Center (NAIC), at the Kitt Peak National Observatory (KPNO), the Cerro Tololo Interamerican Observatory (CTIO) and at the Palomar Observatory (PO), NAIC is operated by Cornell University, KPNO and CTIO by Associated Universities for Research in Astronomy, all under cooperative agreements with the National Science Foundation. The Hale telescope at the PO is operated by the California Institute of Technology under a cooperative agreement with Cornell University and the Jet Propulsion Laboratory. This research was supported by NSF grants AST94-20505 and AST96-17069 to RG, AST95-28860 to MH and by Fondecyt grant 1970735 to LEC.

REFERENCES

- Abell, G., Corwin, H.G. & Olowin, R.P. 1989, ApJS 70, 1
- Bershady, M.A. 1997, in *Dark Matter 1996*, eds. P. Salucci & M. Persic, ASPCS 117, 537 (San Francisco)
- Dale, D.A. 1998, Ph. D. Thesis, Cornell University
- Dale, D.A., Giovanelli, R., Haynes, M.P., Hardy, E. & Campusano, L. 1997, AJ 114, 455
- Dale, D.A., Giovanelli, R., Haynes, M.P., Hardy, E. & Campusano, L. 1998, AJ 115, 418
- Dale, D.A., Giovanelli, R., Haynes, M.P., Campusano, L., Hardy, E. & Borgani, S. 1999, ApJ 510, L11
- Giovanelli, R., Haynes, M.P., Herter, T., Vogt, N.P., da Costa, L.N., Freudling, W., Salzer, J.J. and Wegner, G. 1997a, AJ 113, 22 (G97a)
- Giovanelli, R., Haynes, M.P., Herter, T., Vogt, N.P., da Costa, L.N., Freudling, W., Salzer, J.J. and Wegner, G. 1997b, AJ 113, 53 (G97b)
- Giovanelli, R., Haynes, M.P., Salzer, J.J., Wegner, G., da Costa, L.N. and Freudling, W. 1998a, AJ 116, 2632
- Giovanelli, R., Haynes, M.P., Freudling, W., da Costa, L.N., Salzer, J.J. and Wegner, G. 1998b, ApJ 505, L91

- Postman, M. 1997, in *The Extragalactic Distance Scale*, M. Livio, M. Donahue & N. Panagia, eds., Cambridge:Cambridge, p. 49
- Rix, H.-W., Guhathakurta, P., Colless, M. & Ing, K. 1997, MNRAS 285, 779
- Sandage, A. & Hardy, E. 1973, ApJ 183, 743
- Simard, L. & Pritchett, C.J. 1998, ApJ 505, 96
- Tully, R.B. and Fisher, J.R. 1977, A&A 54, 661
- Vogt, N.P., et al. 1997, ApJ 479, L121
- Zehavi, I., Riess, A.G., Kirshner, R.P. & Dekel, A. 1998, ApJ 503, 483

Table 1. Cluster Positions and Velocities

Cluster	RA(1950)	Dec (1950)	cz_{cmb}	V_{pec}	n
SCI					
N383	010430.0	+321200	4865 ± 32	-6 ± 170	21
N507	012000.0	+330400	4808 ± 99	94 ± 204	14
A262	014950.0	+355440	4664 ± 80	70 ± 133	31
A400	025500.0	+055000	6934 ± 75	-126 ± 227	25
Eridanus	033000.0	-213000	1534 ± 30	-304 ± 74	34
Fornax	033634.0	-353642	1321 ± 45	-109 ± 60	39
Cancer	081730.0	+211400	4939 ± 80	61 ± 172	26
Antlia	102745.0	-350411	3120 ± 100	185 ± 109	27
Hydra	103427.7	-271626	4075 ± 50	-320 ± 142	25
N3557	110735.0	-371600	3318 ± 57	199 ± 155	11
A1367	114154.0	+200700	6735 ± 88	62 ± 191	35
Ursa Major	115400.0	+485300	1101 ± 40	-425 ± 56	30
Cen30	124606.0	-410200	3322 ± 150	310 ± 98	38
A1656	125724.0	+281500	7185 ± 68	212 ± 210	41
ESO508	130954.0	-230854	3210 ± 100	417 ± 128	17
A3574	134606.0	-300900	4817 ± 30	-26 ± 174	20
A2197†	162630.0	+410100	9162 ± 100	-204 ± 384	25
Pavo II	184200.0	-632000	4444 ± 70	137 ± 163	18
Pavo	201300.0	-710000	4055 ± 100	80 ± 219	10
MDL59	220018.0	-321400	2317 ± 75	-503 ± 120	23
Pegasus	231742.6	+075557	3519 ± 80	-186 ± 180	17
A2634	233554.9	+264419	8895 ± 79	-136 ± 270	26
A2666	234824.0	+264824	7776 ± 84	-156 ± 459	9
SCII					
A2806	003754	-562600	7867 ± 80	464 ± 382	10
A114	005112	-215800	17144 ± 143	-578 ± 1111	9
A119	005348	-013200	13141 ± 85	-275 ± 988	6
A2877	010736	-461000	6974 ± 58	-104 ± 489	7
A2877b	010736	-461000	9040 ± 48	307 ± 634	5
A160	011012	+151500	12072 ± 141	280 ± 977	6
A168	011236	-000100	13049 ± 58	679 ± 725	9
A194	012300	-014600	5037 ± 37	-216 ± 302	13
A260	014900	+325500	10664 ± 111	-1175 ± 835	9
A397	025412	+154500	9594 ± 78	553 ± 630	14
A3193	035654	-522900	10522 ± 112	450 ± 668	6
A3266‡	043030	-613500	17782 ± 61	-2700 ± 2345	2
A496	043118	-132100	9809 ± 59	566 ± 513	9
A3381‡	060806	-333500	11510 ± 48	798 ± 868	4
A3407	070342	-490000	12861 ± 136	-179 ± 1235	8
A569	070524	+484200	6011 ± 43	-157 ± 280	13
A634	081030	+581200	7922 ± 42	-222 ± 469	8
A671	082524	+303500	15307 ± 194	-120 ± 838	9
A754‡	090624	-092600	16599 ± 82	-92 ± 3294	3
A779	091648	+335900	7211 ± 101	-100 ± 320	14
A957	101124	-004000	13819 ± 120	-866 ± 974	6
A1139	105530	+014600	12216 ± 71	694 ± 629	11
A1177	110648	+215800	10079 ± 81	51 ± 689	6
A1213	111348	+293200	14304 ± 90	744 ± 899	6
A1228	111848	+343600	10794 ± 34	-603 ± 517	13
A1314	113206	+491900	9970 ± 154	-134 ± 582	8
A3528‡	125136	-284500	16770 ± 139	-1441 ± 1703	3
A1736	132406	-265100	10690 ± 50	-49 ± 887	6
A1736b‡	132406	-265100	14017 ± 84	186 ± 1121	4
A3558	132506	-311400	14626 ± 44	678 ± 981	8
A3566	133606	-351800	15636 ± 87	236 ± 837	9
A3581‡	140436	-264700	7122 ± 126	-139 ± 659	4
A1983b	144724	+170600	11524 ± 62	1291 ± 589	8
A1983	145024	+165700	13715 ± 45	429 ± 1165	7
A2022	150212	+283700	17412 ± 72	-1134 ± 1067	8
A2040	151018	+073700	13616 ± 61	212 ± 839	10
A2063	152036	+084900	10605 ± 53	680 ± 398	18

Table 1—Continued

Cluster	RA(1950)	Dec (1950)	cz_{cmb}	V_{pec}	n
A2147	160000	+160200	10588 ± 85	303 ± 427	19
A2151	160300	+175300	11093 ± 59	312 ± 424	22
A2256	170636	+784700	17401 ± 132	56 ± 998	8
A2295b [†]	175900	+691600	18633 ± 82	-408 ± 1587	4
A2295	180018	+691300	24554 ± 199	-1145 ± 1448	10
A3656	195712	−384000	5586 ± 64	-72 ± 375	6
A3667 [‡]	200830	−565800	16477 ± 94	-3034 ± 1582	4
A3716	204754	−525400	13618 ± 64	359 ± 581	14
A3744	210418	−254100	11123 ± 89	-150 ± 578	11
A2457	223312	+011300	17280 ± 110	-144 ± 946	9
A2572	231554	+182800	11495 ± 100	436 ± 803	5
A2589	232130	+163300	11925 ± 95	-194 ± 804	6
A2593	232200	+142200	12049 ± 86	-761 ± 605	12
A2657	234218	+085200	11662 ± 137	32 ± 844	5
A4038	234506	−282500	8713 ± 63	68 ± 534	7

[†]Includes A2197, A2199.

[‡]Excluded from statistical analysis.